

Math 175: Combinatorics
Homework 3
Due in Discussion Thursday, February 8.

This week we started by finishing our discussion of the birthday problem. Specifically, we talked about the probability of a shared birthday among k people when there are n possible birthdays, but probability of being born on each day is not necessarily uniform. We discussed the case of 2 people and 2 birthdays in detail.

We then talked about proof by induction, which is the subject of Section 2.1 of LVP. We gave some examples including the sum of the first n powers of 2 and the sum of the first n positive integers.

We proved a basic proposition about binomial coefficients, equation (3.2) in Section 3.6 of LVP. We then talked about Pascal's Triangle (LVP Section 3.5) and then many patterns within it.

We also introduced the Fibonacci numbers (LVP Chapter 4) and explained how they arise as sums of shallow diagonals in Pascal's triangle. We gave a formula for the Fibonacci numbers (Theorem 4.3.1 in LVP).

At this point, you should have read all of Chapter 1 of LVP. You should read Section 2.1 on Induction, and it would also be good to read the rest of Chapter 2 on combinatorial tools. We will soon prove the Binomial Theorem, which is covered in Section 3.1. I would recommend reading all of Chapter 3 (except Sections 3.7 and 3.8) and all of Chapter 4. Much of the material on binomial coefficients is also covered in Section 2.2 of the HHM book.

Problems

1. We saw in lecture that if there are two possible birthdays A and B that occur with probabilities p and $1 - p$, the probability that two people have a shared birthday is $2p(1 - p)$. This is a function of a single variable p that reaches a maximum at $p = \frac{1}{2}$. We can see this by taking the derivative with respect to p to find a critical point and checking that it gives a maximum. Suppose there are three possible birthdays: A, B, C and they occur with probabilities p_1, p_2 , and p_3 , respectively. A special case of what we saw in class is that the probability that two people do not share a birthday is given by

$$f(p_1, p_2, p_3) = 2(p_1p_2 + p_1p_3 + p_2p_3).$$

- (a) Suppose that p_3 is fixed. We see that $p_1 + p_2 = 1 - p_3$, so $p_2 = (1 - p_3) - p_1$. Find the value of p_1 that maximizes the function of a single variable

$$f(p_1) = 2(p_1((1 - p_3) - p_1) + p_1p_3 + ((1 - p_3) - p_1)p_3).$$

- (b) When there are n possible birthdays with probabilities p_1, p_2, \dots, p_n , the probability that two people do not have a shared birthday is given by

$$f(p_1, \dots, p_n) = 2 \sum_{1 \leq i < j \leq n} p_i p_j.$$

Suppose that p_2, \dots, p_n are fixed. As above, we can express p_1 in terms of p_2 and these fixed constants. Find the value of p_1 that maximizes the function of a single variable

$$f(p_1) = 2 \sum_{1 \leq i < j \leq n} p_i p_j.$$

- (c) Using the previous result, show that the uniform distribution, where each $p_i = \frac{1}{n}$ maximizes the probability that our two people do not share the same birthday. [Hint: argue by contradiction.]

2. **Exercise 2.5.1 of LVP:** What is the following sum?

$$\frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \frac{1}{3 \cdot 4} + \dots + \frac{1}{(n-1) \cdot n}.$$

Experiment, conjecture the value, and then prove it by induction.

3. **Exercise 2.5.2 of LVP:** What is the following sum?

$$0 \cdot \binom{n}{0} + 1 \cdot \binom{n}{1} + 2 \cdot \binom{n}{2} + \dots + (n-1) \binom{n}{n-1} + n \cdot \binom{n}{n}.$$

Experiment, conjecture the value, and then prove it by induction. (You may want to try to prove this result by induction and also by combinatorial arguments.)

4. In order to get a feel for how induction arguments work, it is instructive to see a bunch of convincing looking induction arguments that are actually wrong. Read Exercises 2.1.12 and 2.1.13 in the LVP book. There are solutions for both of these in the back of the book.

Identify and explain the flaw in the following two induction ‘proofs’.

- (a) CLAIM: For all positive integers n ,

$$\sum_{j=1}^n j = \frac{1}{2} \left(n + \frac{1}{2} \right)^2.$$

We check that this holds for $n = 1$. We suppose that this is true for n and prove it for $n + 1$. We have

$$\begin{aligned} \sum_{j=1}^{n+1} j &= \sum_{j=1}^n j + (n+1) \\ &= \frac{1}{2} \left(n + \frac{1}{2} \right)^2 + (n+1) \text{ by the induction hypothesis} \\ &= \frac{1}{2} \left(n^2 + n + \frac{1}{4} + 2n + 2 \right) \\ &= \frac{1}{2} \left((n+1) + \frac{1}{2} \right)^2. \end{aligned}$$

This is what we wanted to show. Therefore this statement holds for all positive integers n .

- (b) Claim: All positive integers are equal. To prove this claim, we will prove by induction that for all positive integers n the following statement holds.

For any positive integers x and y , if $\max(x, y) = n$, then $x = y$.

(Here $\max(x, y)$ denote the larger of the two numbers x and y , or the common value if both are equal.) We call this statement $P(n)$. Clearly $P(1)$ is true, since $\max(x, y) = 1$ forces $x = 1$ and $y = 1$, so $x = y$.

Suppose that $P(n)$ is true. We will prove that $P(n + 1)$ is true. Let x, y be positive integers such that $\max(x, y) = n + 1$. Then

$$\max(x - 1, y - 1) = \max(x, y) - 1 = n + 1 - 1 = n.$$

By the induction hypothesis, it follows that $x - 1 = y - 1$, and therefore $x = y$. This proves $P(n + 1)$ holds. We now conclude that $P(n)$ holds for all positive integers n , so $\max(1, n) = n$ for any positive integer n . Therefore $1 = n$ for any positive integer n .

5. In class we saw that the sum of the first n positive integers is $\frac{n(n+1)}{2}$. Exercise 2.1.8 in LVP, which has a solution in the back, shows that $\sum_{j=1}^n j^2 = \frac{n(n+1)(2n+1)}{6}$.

- (a) Prove by induction that

$$\sum_{j=1}^n j^3 = \left(\sum_{j=1}^n j \right)^2 = \frac{n^2(n+1)^2}{4}.$$

(b) Prove using induction that

$$\sum_{j=1}^n (-1)^{j-1} j^2 = (-1)^{n-1} \binom{n(n+1)}{2}.$$

6. (a) Use the Hockey-Stick Identity from lecture, equation (3.5) on page 53 of the LVP book, to prove the following:

$$\binom{r}{r} + \binom{r+1}{r} + \cdots + \binom{n}{r} = \binom{n+1}{r+1}.$$

- (b) In lecture we gave a combinatorial proof of the Hockey-Stick identity. Give another. It's fine to give a combinatorial proof of the identity from part (a) of this problem, rather than of the Hockey-Stick result directly. [Hint: Imagine picking $r+1$ items from the set $\{1, 2, \dots, n+1\}$. Classify the ways of doing this according to the value of the largest of those items. Then count in two ways! Also see Exercise 3.6.4 and its Hint in the back of the LVP book.]
7. For this problem, imagine we are flipping a fair coin. That is, on each flip there is a 50% chance of getting H and a 50% chance of getting T. We have already seen that there are 2^n total outcomes when flipping n times, since this is the same as a word of length n made up of H's and T's. These questions look like they are about probability, but they can be interpreted as questions about counting words from a two-letter alphabet with certain properties.
- (a) Flip a coin $2n$ times. What is the probability of getting either $n-1$ heads or n heads?
- (b) Suppose you flip a coin $2n+1$ times. What is the probability of getting either n heads or $n+1$ heads?
- (c) Show that your answers to parts (a) and (b) are equal. You can do this using a binomial identity from lecture, or (and this solution is nicer), you can give a combinatorial interpretation of what's going on here. [Hint for the combinatorial proof: ask yourself what is the number of heads you must get in the first $2n$ flips and the corresponding number of heads in the next flip in order to get either n or $n+1$ heads in $2n+1$ flips.]